SCIENCE LIBRARY

MAR 24 1922

VOL. 49 JANUARY, 1922 No. 1

BULLETIN

OF THE

TORREY BOTANICAL CLUB

Editor

ALEXANDER WILLIAM EVANS

Associate Editors

JEAN BROADHURST
JAMES ARTHUR HARRIS
GEORGE T. HASTINGS
MARSHALL AVERY HOWE
MICHAEL LEVINE
GEORGE ELWOOD NICHOLS
ARLOW BURDETTE STOUT
NORMAN TAYLOR

CONTENTS

Epiphyllous plants of certain regions in Jamaica (Plate 1)
LOUIS J. PESSIN 1

The Genus Catinella . . . . . ELIAS J. DURAND 13

INDEX TO AMERICAN BOTANICAL LITERATURE 23

PUBLISHED FOR THE CLUB

PRESS OF
THE NEW ERA PRINTING COMPANY
LANCASTER, PA.

THE NEW ERA PRINTING COMPANY

LANCASTER, PA.

Is prepared to execute in first-class and satisfactory manner all kinds of Printing, Electrotyping and Binding. Particular attention given to the work of Schools, Colleges, Universities and Public Institutions

Books, Periodicals

Technical and Scientific Publications

Monographs, Theses, Catalogues

Announcements, Reports, etc.

All Kinds of Commercial Work

Publishers will find our product ranking with the best in workmanship and material, at satisfactory prices. Correspondence solicited. Estimates furnished.

THE NEW ERA PRINTING COMPANY

American Fern Journal

An illustrated quarterly devoted to the general study of ferns. Subscription, with membership in the AMERICAN FERN SOCIETY, $1.50 per annum; without membership, $1.25; to foreign subscribers, $1.35; life membership in the Society, $25.00. Volume I, six numbers, $2.00; other volumes, $1.00 each. Matter or publication should be sent to R. C. BENEDICT, 322 East 19th Street, Brooklyn, N. Y.; subscriptions and other business communications to E. J. WINSLOW, Auburndale, Mass.

LABORATORY SUPPLIES

OF EVERY DESCRIPTION

MICROSCOPES AND ACCESSORIES

Slides, Coverglasses, Stains, Glassware, Etc.

A well assorted stock always on hand

MEYER CAMERA & INSTRUMENT CO., Inc.

89 Third Avenue **NEW YORK**

Please mention the "Bulletin" when answering advertisements.

Gen.
Wash

CONTENTS

PESSIN, LOUIS J. Epiphyllous plants of certain regions in Jamaica (plate 1) 1
DURAND, ELIAS J. The genus Catinella 15
MUNZ, PHILIP A., and JOHNSTON, IVAN M. Miscellaneous notes on plants of Southern California—I 31
FARR, CLIPPORD H. Quadripartition by furrowing in Sisyrinchium (plate 2) 51
ROUND, EDA M. A modern plant fossil 63
HAZEN, TRACY E. The phylogeny of the genus Brachiomonas (plates 3, 4) 75
STEVENS, O. A. New records and other notes on North Dakota plants. 93
YUNCKER, T. G. Three new species of Cuscuta from Mexico 107
HAZEN, TRACY E. New British and American species of Lobomonas: a study in morphogenesis of motile algae (plates 5, 6) 123
ANDERSON, FLORA. The development of the flower and embryogeny of Martynia louisiana (plates 7, 8) 141
OVERHOLTS, L. O. Mycological notes for 1920 (plate 9) 163
GRAFF, PAUL W. Unreported plants from Glacier National Park 175
OSTERHOUT, GEO. E. Two new plants from Colorado 183
ARTHUR, JOSEPH CHARLES. New species of Uredineae—XIV 189
SMITH, CHARLES PIPER. Studies in the genus Lupinus—VII. L. succulentus and L. niveus 197
HOWE, MARSHALL A., and HOLLICK, ARTHUR. A new American fossil hepatic 207
COCKERELL, T. D. A. A new genus of fossil Liliaceae 211
GRAFF, PAUL W. Philippine Basidiomycetes—V 223
DOSDALL, LOUISE. Occurrence of the pyncial stage of Puccinia Taraxaci 235
BARTRAM, EDWIN B. Midwinter botanizing in southern Arizona 237
BRITTON, N. L., and ROSE, J. N. Two new genera of Cactaceae 251
RUSBY, H. H. New species of trees of medical interest from Bolivia ... 259
ASHE, W. W. Notes on trees and shrubs of the southeastern United States 265
THARP, B. C. Commelinantia, a new genus of Commelinaceae (plates 10, 11) 269
ROWLEE, W. W. The genus Costus in Central America (plates 12–15) .. 283
SHIMEK, B. Quercus lyrata in Iowa (plates 16, 17) 293
PORTERPIELD, W. M., JR. References to the algae in the Chinese classics. 297
DODGE, B. O. A Lachnea with a botryose conidial stage 301

DORAN, W. L. Effect of external and internal factors on the germination of fungous spores 313
MUNZ, PHILIP A., and JOHNSTON, IVAN M. Miscellaneous notes on plants of Southern California—II 349
MACKENZIE, KENNETH KENT. Notes on Carex—XII 361
OLDENBUSH, CARRIE. Stimulation of plants by carbon disulphide ... 375

INDEX TO AMERICAN BOTANICAL LITERATURE, 23, 45, 65, 111, 159, 185, 215, 253, 277, 307, 341, 391

INDEX TO VOLUME 49 399

Dates of Publications

No. 1, for January.	Pages 1– 30.	Issued March 21, 1922.
No. 2, for February.	31– 50.	April 4, 1922.
No. 3, for March.	51– 74.	April 12, 1922.
No. 4, for April.	75–122.	May 18, 1922.
No. 5, for May.	123–162.	June 15, 1922.
No. 6, for June.	163–188.	July 8, 1922.
No. 7, for July.	189–222.	August 10, 1922.
No. 8, for August.	223–258.	August 31, 1922.
No. 9, for September.	259–282.	September 22, 1922.
No. 10, for October.	283–312.	October 11, 1922.
No. 11, for November.	313–348.	November 9, 1922.
No. 12, for December.	349–408.	January 27, 1923.

Errata

Page 2, line 8, for "*arboreum*" read "*arborum.*"
Page 6, footnote, for "*Caraguta*" read "*Caraguata.*"
Page 14, line 7, for "PLATE 5" read "PLATE 1."
Page 14, line 2 from bottom, for "*Caraguta*" read "*Caraguata.*"
Page 32, line 11 from bottom, for "*muticus*" read "*mutica.*"
Page 38, line 13 from bottom, for "*racemosa*" read "*racemosum.*"
Page 76, line 1, for "gareement" read "agreement."
Page 77, line 12 from bottom, for "genera" read "general."
Page 92, line 4, for "ertracted" read "retracted."
Page 100, line 4, for "AMSINKIA" read "AMSINCKIA."
Page 101, line 13, for "SIBERICA" read "SIBIRICA."
Page 109, line 6, for "wartius" read "Martius."
Page 109, line 10 from bottom, for "falsl" read "false."
Page 125, line 7 from bottom, insert the word "are" after "sure."
Page 127, line 16, insert a semicolon after "posterioribus."

Errata (*Continued*)

Page 127, line 23, for "zsooporis" read "zoosporis."
Page 143, Fig. 7, the lower right hand "S" should be "S[1]."
Page 156, line 12, for "1902" read "1901."
Page 166, lines 1 and 6 from bottom, for "*hyalina*" read "*hyalinus.*"
Page 167, line 14, for "HYALINA" read "HYALINUS."
Page 180, line 12, for "PENTSTEMON" read "PENSTEMON."
Page 189, line 9, for "*Pentstemon*" read "*Penstemon.*"
Page 195, line 5, for "*Bermudianum*" read "*Bermudiana.*"
Page 197, line 8 from bottom, omit "with."
Page 208, after explanation of FIG. 1 add "× 3."
Page 224, line 16, for "*corrugatus*" read "*corrugata.*"
Page 226, line 9, for "*Polystitictus*" read "*Polystictus.*"
Page 228, line 18, for "MULITPLEX" read "MULTIPLEX."
Page 241, line 16, for "NOTHOLEANA" read "NOTHOLAENA."
Page 242, line 1, for "MACHROSTACHYA" read "MACROSTACHYA."
Page 242, line 16, for "RINGENS" read "RIGENS."
Page 244, line 8 from bottom, for "VAUQUELINA" read "VAUQUELINIA."
Page 245, lines 24 and 26, for "PAROSELLA" read "PAROSELA."
Page 247, line 7, for "CHOLOROTICA" read "CHLOROTICA."
Page 248, line 12 from bottom, for "PENTSTEMON" read "PENSTEMON."
Page 271, line 22, for "spathe" read "sheath of leaf."
Page 271, line 6 from bottom, for "leaves" read "branches."
Page 275, line 4, for "*Tinantia*" read "*Tradescantia.*"
Page 291, line 15 from bottom, for "herbarium of Cornell University" read "U. S. National Herbarium."
Page 292, line 19, for "Aented" read "Oersted."
Page 320, lines 5 and 13 from bottom, for "*Entothia*" read "*Endothia.*"
Page 323, line 7, for "*Plasmopora*" read "*Plasmopara.*"

Vol. 49 No. 1

BULLETIN

OF THE

TORREY BOTANICAL CLUB

JANUARY, 1922

Epiphyllous plants of certain regions in Jamaica*

LOUIS J. PESSIN

(WITH PLATE I AND ONE TEXT FIGURE)

The great variety and luxuriant growth of the epiphyllous plants seen in Jamaica in the summer of 1919 led the writer to study them from two points of view: first, their distribution in relation to climatic conditions; second, their histological relation to the leaves bearing them, in order to determine whether these epiphytes are to any degree parasitic. The study of the distribution and environment of the epiphyllous plants was made in the field. The relation of the epiphyll to its supporting leaf was studied by histological methods in the laboratory.

HISTORICAL

Very little is known about epiphyllous plants today, in spite of the fact that they have long interested botanists. As early as 1875 Berkeley (1) reported the occurrence of epiphyllous lichens growing on the leaves of the tea plant; some of these apparently epiphyllous forms showed marked parasitic adaptation. Such forms were found in abundance in warm moist regions. Two of the forms observed by Berkeley were regarded as species belonging to the genus *Strigula*.

A few years after Berkeley's paper appeared, Cunningham (2) reported on an epiphyllous lichen and an epiphyllous alga, *Mycoidea parasitica*, parasitic on leaves of *Mangifera indica*, *Croton* and *Rhododendron*.

[The BULLETIN for December (48 : 315–363) was issued February 28, 1922.]

* Botanical Contribution No. 69 from the Johns Hopkins University.

Marshall Ward (6) worked on an epiphyllous lichen which he believed to be the one worked on by Cunningham. He observed no penetration of the tissues of the supporting leaf by the lichen and therefore decided that it was not parasitic. He did find, however, one or more cross walls in the palisade cells of the supporting leaf beneath the epiphylls, which formed three or four layers of these cells instead of the normally single layer.

Schmidle (5) found species of *Trentepohlia arboreum* and *T. aurea* as well as *Phycopeltis microcystis* quite common as epiphyllous forms in Samoa. Massart (3) found that some orchids pass their entire existence on a single leaf. He found also that the epiphyllous floras differ in different regions. Algae and liverworts are the most common epiphyllous forms, while mosses acquire the epiphyllous habit only in very humid ravines.

OBSERVATIONS AND METHODS

Specimens of epiphyllous plants were collected in various regions in the Blue Mountains of Jamaica. The field observations were made during June and July, 1919, in the following places: Cinchona Plantation (5,000 feet), Morce's Gap (4,934 feet), New Haven Gap (5,600 feet), Green River Valley (3,000 feet), Markham Hill (5,800 feet), Vinegar Hill Road (4,500 feet), Clyde River Valley (4,500 feet) and Blue Mountain Peak (7,428 feet).

Material was either pressed or preserved in glycerine in 4 per cent formaldehyde, or in 60 per cent alcohol. For careful histological study, material was fixed in chromoacetic acid and preserved in 70 per cent alcohol until finally dehydrated, embedded, cut and stained with the Flemming's triple stain. In some cases material was left unstained for comparison.

The Cinchona Plantation is situated on a spur running south from the Blue Mountains at five thousand feet above sea level. The temperature varies between the extremes of 48° and 82° F. The vegetation of the garden about the residence of the Cinchona Plantation consists largely of introduced plants, many of which are Himalayan, Australian and Cape of Good Hope species. There are also native tree ferns, orchids, bromeliads, and many other transplanted native species. The banks along the trails outside the garden are covered mostly by native ferns interspersed

with, or on the north side of Cinchona Hill largely replaced by, *Hedychium coronarium*.

During the wet season Cinchona is enveloped in clouds for days at a stretch and the rains are very heavy and frequent. In spite of this abundance of moisture at certain times of the year, epiphyllous plants are rather scarce on the south side of Cinchona Hill. Only one hepatic, two species of algae and a few kinds of lichens were found growing there. Of these *Trentepohlia* grows in damp or shaded places on the ground, on the bark of trees and on the leaves of certain plants. Lichens, the most common of which belong to the genus *Sticta*, grow abundantly on certain plants. Some of the leaves of *Callistemon lanceolatum*, for instance, are almost completely covered by this lichen. In a few cases the leaves of *Hedychium* were found sprinkled with a reddish disk-like alga belonging to the genus *Phycopeltis*. The epiphyllous flora of this region is evidently not a rich one (see TABLE I, A).

At Morce's Gap, two miles north of Cinchona and at about the same level, is found a characteristic tropical montane rain forest. Here moisture is abundant at all seasons of the year. On the windward side of the mountains everything reeks with moisture; clouds drift constantly through these dense forests and rain falls frequently during eight months of the year. The dense vegetation, except on the ridges, is usually dripping wet. Very rarely do the sun's rays penetrate to the floor of these forests. Ferns carpet the floor and cover the walls of the steep-sided ravines. Tree ferns, such as *Alsophila* and *Cyathea*, are very common, many of them having trunks four or five inches thick and thirty feet high or more. Species of the genera *Marattia*, *Danaea*, *Polypodium* and *Elaphoglossum* are very abundant. Such forms as *Blechnum attenuatum* and many species of *Trichomanes*, *Hymenophyllum*, *Polypodium* and *Elaphoglossum* have been compelled to adopt either the climbing or the epiphytic habit in the intense competition for sunlight. *Podocarpus*, *Hedyosmum*, species of *Piper* and many other dicotyledonous trees and shrubs, together with numerous herbaceous plants, are scattered through these woods, increasing the struggle for existence and making the competition for light more keen.

In such an environment epiphyllous plants are very abundant.

TABLE I

EPIPHYLLOUS PLANTS OF JAMAICA

Region	Supporting plant	Algae	Lichens	Hepaticae	Musci
A. Cinchona Plantation, 5000 ft. alt.; south exposure.	*Callistemon lanceolatum*		*Parmelia* sp. *Sticta* sp. *Usnea* sp. A form of Pilocarpaceae	*Cololejeunea diaphana*	
	Hedychium coronarium	*Phycopeltis* sp. *Trentepohlia aurea*	*Lopadium* sp.		
	Pittosporum tobira	*Phycopeltis* sp.			
B. Morce's Gap, 4934 ft. alt.; north exposure.	*Elaphoglossum latifolium*	*Desmidium* sp. *Nostoc* sp. *Phycopeltis* sp. *Pinnularia* sp. *Scytonema* sp. *Staurastrum* sp.		*Cololejeunea* sp. *Cololejeunea diaphana* *Crossotolejeunea* sp. *Diplasiolejeunea pellucida* *Metzgeria furcata* *Microlejeunea* sp. (?)	
	Hedychium coronarium	*Phycopeltis* sp.		*Crossotolejeunea* sp. *Aphanolejeunea* sp. *Metzgeria furcata* *Cololejeunea diaphana* *Diplasiolejeunea pellucida* Schiffn.	
	Hedyosmum arborescens Sw.			*Crossotolejeunea* sp. *Lejeunea flava*	

TABLE I—*Continued*

Region	Supporting plant	Algae	Lichens	Hepaticae	Musci

	Hedyosmum arbor-escens Sw.			*Lejeunea flava*	

TABLE I—*Continued*

Region	Supporting plant	Algae	Lichens	Hepaticae	Musci
B. Morce's Gap—*Continued.*	*Mangifera indica*	*Phycopeltis* sp.	Undetermined species*		
	Podocarpus coriaceus	*Phycopeltis* sp.		*Crossotolejeunea* sp. *Lejeunea flava*	
	Polypodium sp.		Undetermined species*		*Clastobryum americanum*
C. New Haven Gap, 5600 ft. alt.; north exposure.	*Stelis ophioglossoides*	*Phycopeltis* sp. *Trentepohlia aurea*			
	Trichomanes sp.			*Crossotolejeunea* sp.	
	Blechnum attenuatum	*Phycopeltis* sp. *Pinnularia* sp. *Nostoc* sp. *Scytonema* sp.		*Cololejeunea diaphana* *Crossotolejeunea* sp. *Diplasiolejeunea pellucida* *Drepanolejeunea campanulata* *Lophocolea* sp. *Metzgeria furcata* *Taxilejeunea* sp.	*Miltenothamnium reptans*
	Clusia havetioides	*Pinnularia* sp. *Nostoc* sp. *Scytonema* sp.	*Parmelia* sp.	*Aphanolejeunea* sp. *Aphanolejeunea sicaefolia* *Diplasiolejeunea pellucida* *Drepanolejeunea campanulata* *Cololejeunea diaphana* *Lejeunea flava* *Metzgeria furcata* *Microlejeunea* sp.	*Miltenothamnium reptans*

* Up to the present the writer has been unable to have some of the epiphyllous lichens definitely determined.

TABLE I—*Continued*

Region	Supporting plant	Algae	Lichens	Hepaticae	Musci
New Haven Gap—*Continued.*	*Monoclea Gottschei*				*Mittenothamnium reptans*
	Plagiochila sp.			*Riccardia* sp.	*Mittenothamnium reptans*
	Rhododendron sp.		*Phylloporina epiphylla* Undetermined species		
D. Vinegar Hill Road, 4500 ft. alt.: north and west exposures.	*Asplenium* sp.			*Cololejeunea* sp. *Cololejeunea diaphana* *Crossotolejeunea* sp. *Diplasiolejeunea pellucida* *Drepanolejeunea campanulata* *Metzgeria furcata*	
	Clusia havetioides	*Nostoc* sp. *Phycopeltis* sp. *Pinnularia* sp.	*Parmelia*	*Cololejeunea* sp. *Diplasiolejeunea pellucida* *Drepanolejeunea campanulata* *Lejeunea flava* *Lophocolea* sp. *Metzgeria furcata* *Taxilejeunea* sp.	
	*Elaphoglossum latifolium**	*Chroococcus* sp. *Gleocapsa* sp. *Nostoc* sp. *Phycopeltis* sp. *Scytonema* sp.		*Cololejeunea diaphana* *Crossotolejeunea* sp. *Lejeunea flava* *Metzgeria furcata*	*Mittenothamnium reptans*

* On a leaf of this species the seed plant *Caragata Sintenesii* was likewise found living as an epiphyll.

* On a leaf of this species the seed plant *Caraguta Sintenesii* was likewise found living as an epiphyll.

TABLE I—*Continued.*

Region	Supporting plant	Algae	Lichens	Hepaticae	Musci
D. Vinegar Hill Road—*Continued.*	*Epidendrum* sp.		Undetermined species.	*Drepanolejeunea campanulata*	
	Polypodium Phyllitidis			*Cololejeunea* sp. *Cololejeunea diaphana* *Crossotolejeunea* sp. *Diplasiolejeunea pellucida* *Drepanolejeunea campanulata* *Drepanolejeunea crucianella* *Lophocolea* sp. *Metzgeria furcata* *Taxilejeunea* sp.	*Fissidens* sp.
E. Blue Mountain Peak, 7428 ft. alt.; northeast exposure.	*Podocarpus coriaceus*	*Phycopeltis* sp.			
F. Clyde River Valley, 4000 ft. alt.; north and west exposures.	*Citrus Limonium*	*Phycopeltis* sp. *Scytonema* sp. *Trentepohlia aurea*	*Coenogonium* sp. *Collema* sp. *Lecanora* sp. *Pannaria rubiginosa* *Pyrenula* sp. Undetermined species.	*Aphanolejeunea* sp. *Cololejeunea* sp. *Diplasiolejeunea pellucida* *Drepanolejeunea campanulata* *Lejeunea flava* *Metzgeria furcata*	

Here one finds plants of many types growing on the soil; on these grow various plants as epiphytes and on these in turn grow other epiphytes. A remarkable diversity of epiphytic forms can be found on one plant, and some of the forms are wonderfully adapted to their epiphytic mode of life. These epiphytes occur on every patch of dead or living plant surface that receives even a moderate supply of light. One finds on a single leaf here such forms as algae, lichens, hepatics, mosses and, at times, even seed plants, not to mention occasional fungi and minute epiphyllous animals. It is rather strange that not a single fern prothallus has been found on any of the leaves of this region, though numerous sporangia were present and probably spores also on many different kinds of leaves. In TABLE I, B, are listed the different kinds of epiphyllous forms found near Morce's Gap.

The climatic conditions of New Haven Gap do not differ markedly from those of Morce's Gap. The temperature is nearly the same, ranging annually from 40.5° to 83° F., and the floor of the forest is constantly moist and shaded. Here again one finds the characteristic rain forest vegetation. The epiphyllous plants here are quite as abundant as at Morce's Gap and consist mainly of algae, hepatics, and mosses. The leaves of the climbing plants of this locality are well inhabited by epiphyllous forms. The leaves of *Clusia havetioides* and of such ferns as *Blechnum attenuatum*, *Elaphoglossum latijolium* and *Polypodium Phyllitidis* serve most frequently as the substratum for epiphylls (see PLATE 1). A glance at TABLE I, C, gives one a fair notion of the groups of epiphyllous plants found in the New Haven Gap region.

Going northward from Morce's Gap along the upper eastern slopes of the Mabess Valley the trail is particularly interesting as it presents a splendid view of the ravines. The crowns of the tree ferns carpet many side ravines of the valley. The huge lianes, the numerous orchids, the wild pines and the purplish filmy ferns mantle the branches of many trees and the species of *Asplenium*, *Blechnum* and *Polypodium*, along with the walking fern (*Polystichum plaschnichianum*), spread over the banks. One does not have to walk off the trail to find numerous epiphyllous plants. The upper leaves of many of the low trees on the Vinegar Hill Road are completely covered by them. TABLE I, D, shows the great variety of epiphyllous forms found in this region.

On TABLE I, E and F, are indicated the epiphyllous plants observed on the Blue Mountain Peak and in the Clyde River Valley. In the former, though there is always an abundance of moisture, the epiphyllous flora is not a rich one, owing probably to the frequent heavy rains, the rather low temperature and the occasional strong winds. Those of the Clyde River Valley occur chiefly on trees growing along streams, where there is sufficient shade and a humid atmosphere.



FIG. 1. Part of transverse section of a leaf of *Elaphoglossum latifolium* (*E*) with the rhizoids (*Rh*) of an epiphyllous liverwort (L^1) pressed into its cuticle. This liverwort in turn bears another epiphyllous liverwort (L^2), which is likewise attached by its rhizoids (*Rh*).

In general, observations in these Blue Mountains of Jamaica show that the epiphyllous members of each group of plants are limited to certain regions where climatic conditions are suitable for such growth. Thus, Algae, Hepaticae, and Musci usually thrive best on the windward side of the mountains and are abundant only in very damp and very shady woods. Epiphyllous lichens, on the other hand, do not require very warm, humid, and shady places but are also found in relatively dry regions. It was rather difficult to ascertain with any degree of definiteness the exact age of the leaves which bore epiphyllous plants, but it is safe to state that none are present on very young leaves. As a rule they do not appear on leaves until the latter are one or two seasons old.

In order to determine the relation between the epiphyllous forms and the plants on which they grow, cross sections were made of the leaves of *Elaphoglossum latifolium*, *Clusia havetioides*,

Polypodium Phyllitidis and *Mangifera indica*. The leaves of *Elaphoglossum latifolium* were those chiefly used because of the abundance on them of epiphyllous liverworts in various stages of development. Sections of leaves of *Mangifera indica* were also studied to determine the relation of an epiphyllous lichen to the mango leaf.

Sections of the leaves bearing epiphyllous hepatics showed characteristic thickened, club-like rhizoids arising in groups from the stem of the creeping liverwort. These rhizoids are sunken into the cuticle of the supporting leaf as if pressed in by some external force. The rhizoids broaden at their lower ends and flatten out as is shown in TEXT-FIG. 1. This is true even of the rhizoids of one epiphyllous liverwort when growing upon another, for one sometimes finds two or more tiers of epiphylls growing upon a fern or a dicotyledon. The rhizoids of the upper hepatic are in such cases decidedly flattened against the delicate leaves of the lower liverworts. This too is shown in the figure. The important point to be noted is that the rhizoids of the epiphyllous form were not observed to penetrate the tissues of the supporting leaf. Nor was there evident any abnormality in the internal structure of the supporting leaf, such as might conceivably be produced by the shade or by the weight of the epiphylls on the leaf or, possibly, by some substance such as an organic acid excreted by the rhizoids on the liverwort.

The leaves of *Mangifera indica* bore numbers of orange-red disk-like plants of a species of *Phycopeltis*. Among these orange-red specks there were also minute grey patch-like lichens resembling those described by Cunningham (3) and Ward (9). The relation between these lichens and the mango leaf on which they grew has not been definitely determined. The sections of the mango leaf thus far studied showed no modifications of its tissues and no penetration by the lichens.

DISCUSSION

The problem of the evolutionary origin and individual development of the epiphyllous plants is a complex one which still requires much careful investigation. Schimper (4) observed that epiphytic seed plants commonly originate on their support through seeds.

The scattered seeds germinate on the trunk, branches, or leaves of a tree or shrub and, with adequate moisture and other nutritive material, develop into epiphytes. He divides the fruits and seeds of epiphytes into three categories: first, those transported by birds, monkeys and other tree-inhabiting animals; second, those which are very light and may be transported by the wind and which may readily catch in the crevices of the bark and germinate there; third, those seeds which are especially adapted to be carried by the wind. Such seeds usually possess either hairs or wings or inflated coats which aid their distribution.

It is conceivable that a great many of the epiphyllous forms here being considered arise from spores, for spores have been found on many leaves. The epiphyllous liverworts evidently arise often from gemmae, for many such gemmae in the process of development have been found on leaves. The spores of liverworts, on the contrary, have rarely been found germinating on leaves. The observations thus far made indicate that many an epiphyllous hepatic or moss makes its start on the host by creeping up the stem of the support from the ground or up the petiole from the stem. As it climbs upward it dies off from below and when its younger part reaches the leaves it spreads over the surface and becomes strictly an epiphyllous form. Secondly, epiphyllous forms may be lodged upon the host by the agency of water. When during a rain the water flows down over the leaves and branches of a tree in the rain forest, it breaks off the hepatics and mosses and other plants from the surface of the stem or leaf and washes them down on the leaves below, where they continue to grow, covering the leaf in the course of time with a dense mat. Such is probably the explanation of the fact that in most cases the lower leaves possess the greatest number of epiphyllous plants, though, of course, lower leaves are older, better shaded, and are surrounded by moist air. It is probable also that the epiphyllous plants found on the dorsal sides of certain overturned leaves arise in the manner just described. The fact that no epiphyllous plants were observed on the lower sides of leaves still in normal position is evidence that these plants are propagated largely by these fragments. If it were commonly by spores we should expect these, at least occasionally, to stick to the underside of a

leaf. Such cases, however, are extremely rare. In a third possible method of dispersal, spores, gemmae, or seeds may be washed down by the dripping water from above, or may be blown from adjacent plants by the wind. This method is particularly common among algae and lichens. The first two types of vegetative propagation seem to be the commoner ones, for in many instances the epiphyllous forms growing on the stem have actually been observed to spread over the surface of the leaf. Moreover, many of the forms found on the leaves occur likewise on the stem of the supporting plant. This is particularly true of hepatics and mosses. In lichens, however, the case may be different. There the alga, as well as the component fungus, may often be present separately on the leaf; each may reach the leaf in the form of loose spores or fragments, and there combine to give rise to the lichen itself.

Massart (3) failed to find in Java any epiphyllous pteridophytes and phanerogams. In the tabulated lists of epiphyllous plants of Jamaica in this paper it will be noted that no pteridophytes have been found. Although many fern sporangia were observed among the epiphylls by the writer, no germinating fern spores or fern prothallia were seen. The writer, however, found a few epiphyllous phanerogams. These were very young plants of the genus *Caraguata* attached to the surface of the leaf, and apparently doing well. How much further their development would continue is hard to tell, for the substratum is not firm enough to support a mature wild pine; before the epiphyllous wild pine could have time to mature, the supporting leaf would probably be shed. The fact is, at all events, that no mature wild pines were found growing on leaves. According to Massart, the epiphyllous plants do not constitute a similar flora everywhere; in some localities this consists solely of thallophytes, in others of bryophytes. This is also true for Jamaica, where in certain regions only one type occurs, while in other regions numerous diverse forms are present.

Massart's suggestion that the leaf-inhabiting fungus has been derived from the epiphyllous plant is at least favored by the relation to its host of the alga *Cephaleuros*, which may have either the parasitic or the epiphytic mode of life. On the other hand, there are still some fungi which possess the epiphyllous habit, such as *Meliola astrina* and a few others. There seems little doubt that

a careful study of epiphyllous plants would throw some light on the origin of parasitism, such as that of leaf-infesting fungi.

SUMMARY

1. Epiphyllous plants of diverse selected regions in Jamaica were studied in the field and collected for laboratory study in Baltimore.

2. The results of the field observations were tabulated so as to give at a glance a general idea of the distribution of these epiphyllous plants in relation to the altitude, average temperature, and average humidity of their various habitats.

3. The possible modes of attaining the epiphyllous position are:

(*a*) Continual growing forward and dying off below, of hepatics and mosses, until they reach and cover the leaf.

(*b*) By distribution of spores, gemmae or seeds through the agency of rain and wind. This is especially true for algae and fungi.

(*c*) Propagation by displaced fragments of mature lichens, hepatics, and mosses.

4. An abundance of humidity, moderate temperature, and a relatively calm atmosphere encourage an abundant epiphyllous growth.

5. Histological studies of epiphyll-bearing leaves of *Elaphoglossum latifolium*, *Clusia havetioides* and *Mangifera indica* revealed no apparent penetration of the leaf by any part or organ of the epiphyll.

6. The rhizoids of epiphyllous hepatics are especially adapted for the epiphyllous habit by becoming thickened, by flattening out and by being pressed into the cuticle of the leaf, thus firmly attaching the epiphyll to the leaf and preventing its being blown or washed off easily.

7. It is conceivable that the shade produced by the epiphyll, the accumulation of the organic material, the possible excretion of acids and other substances by its rhizoids may have some physiological effect on the supporting leaf.

8. A careful study of the epiphyllous mode of life will, it is believed, bring out some close relations between parasitism and

the epiphyllous habit. Parasitism is possibly an advanced, specialized phase of epiphytism.

The writer is indebted to the Botanical Department of the Johns Hopkins University for the opportunity of going to Jamaica. The writer wishes to thank particularly Professor Duncan S. Johnson for suggesting this problem, for taking the photographs reproduced on PLATE 5, and for helpful advice in carrying on the work. Acknowledgments are due Professor Alexander W. Evans for the determination of most of the epiphyllous hepatics; to Mrs. N. L. Britton of the New York Botanical Garden for the determination of some of the epiphyllous mosses; and to Professors C. C. Plitt and Lincoln W. Riddle for the determination of certain epiphyllous lichens.

LITERATURE CITED

1. **Berkeley, M. J.** On the thread blight of tea. Quart. Jour. Microsc. Sci. **15**: 130–133. *f. 1–4.* 1875.
2. **Cunningham, D. D.** On *Mycoidea parasitica*. Trans. Linn. Soc. London II. **1**: 301–316. *pl. 42, 43.* 1877.
3. **Massart, J.** Les végétaux épiphylles. Ann. Jard. Bot. Buitenzorg. Suppl. 2: 103–108. 1898.
4. **Schimper, A. F. W.** Botanische Mittheilungen aus den Tropen **2**. Die epiphytische Vegetation Americas. Jena. 1888.
5. **Schmidle, W.** Einige Baumalgen aus Samoa. Hedwigia **36**: 277–287. *pl. 6–9.* 1897.
6. **Ward, H. M.** On the structure, development and life history of a tropical epiphyllous lichen. Trans. Linn. Soc. London II. **2**: 87–119. *pl. 18–21.* 1883.

Explanation of plate 1

FIG. 1. Leaf of *Polypodium Phyllitidis*, showing epiphyllous hepatics and mosses.

FIG. 2. Leaf of *Elaphoglossum latifolium*, showing a luxuriant growth of epiphyllous plants; a small wild pine, *Caraguta Sintenesii*, can be seen on the right hand margin.

The genus Catinella

ELIAS J. DURAND

The genus *Catinella* was established by Boudier* for a single species, of rather indefinite relationship, which has commonly been identified as *Peziza olivacea* Batsch. While Batsch's plant is indeterminable when tested according to present day standards of specific limitations, the fungus which has gone by that name is well known, being clearly described by Massee,† and nicely figured by Boudier.‡ The synonymy listed below emphasizes not only the truth of Boudier's statement that the species has been included sometimes in one genus, sometimes in another, but that it has been referred by various writers to several different families as well. The inoperculate asci and intense reaction to caustic potash remove it from the Pezizaceae, where the bright color and fleshy consistency of the fresh young plants at first seem to place it, and to which it has been referred by Saccardo, Rehm and others. The somewhat gelatinous nature of the excipulum has suggested the Bulgariaceae to Fries, Karsten and others, but that character is certainly not conspicuous enough to associate the plant with *Bulgaria inquinans* or *Sarcosoma rufum*. Moreover, a truly gelatinous tissue does not become friable on drying. The present writer is inclined to the opinion of Berkeley, Phillips, Massee and Boudier that its affinities are more properly with the Patellariaceae, such, for example, as *Karschia*. The genus may be characterized as follows:

A genus of the Patellariaceae. Ascomata fleshy and somewhat gelatinous when fresh, becoming friable when dry, sessile, attached to the substratum by radiating, dark fibers; excipulum entirely parenchymatous, becoming deep violet with KOH; asci opening by a pore, not blue with iodine; spores eight, continuous, pale brown; asci and paraphyses agglutinated at the tips to form an epithecium.

---

* Hist. Class. Disc. Eur. 150. 1907.

† Brit. Fungus-Fl. 4: 94. 1895.

‡ Icon. Myc. 3. *pl. 432.*

Species two, as follows:

Spores 7–11 × 4–5 μ.	1. *C. nigro-olivacea.*
Spores 13–15 × 6–7 μ.	2. *C. elastica.*

1. **Catinella nigro-olivacea** (L. v. S.) Durand, comb. nov.

?*Peziza olivacea* Batsch, Elench. Fung. 127. *pl. 12. f. 51.* 1783.
Peziza nigro-olivacea L. v. S. Syn. Car. 121. 1822.
Patellaria pulla β nigro-olivacea Fr. Syst. **2**: 160. 1822.
Bulgaria nigrita Fr. Elench. **2**: 16. 1830.
Lemalis rufo-olivacea L. v. S. Syn. N. Am. n. 1089. 1834.
Rhizina nigro-olivacea Curr. Trans. Linn. Soc. **24**: 494. *pl. 51. f. 10–12* (*fide* Phillips).
Peziza viridi-atra B. & C. Jour. Linn. Soc. Bot. **10**: 369. 1868.
Patellaria violacea B. & Br. Jour. Linn. Soc. Bot. **14**: 108. 1875 (*fide* Massee *l.c.*).
Patellaria hirneola B. & Br. Jour. Linn. Soc. Bot. **14**: 108. 1875 (*fide* Massee *l.c.*).
Patellaria applanata B. & Br. Jour. Linn. Soc. Bot. **14**: 108. 1875.
Peziza fuscocarpa Ell. & Hol. Jour. Myc. **1**: 5. 1885.
Patellaria olivacea Phill. Brit. Disc. 361. 1887 (Batsch ?).
?*Humaria olivacea* Sacc. Syll. **8**: 148. 1889 (Batsch ?).
Pezicula viridi-atra Sacc. Syll. **8**: 315. 1889.
Phaeopezia fuscocarpa Sacc. Syll. **8**: 474. 1889.
Bulgariella pulla β nigro-olivacea Sacc. Syll. **8**: 638. 1889.
Bulgariella nigrita Sacc. Syll. **8**: 638. 1889.
Patinella violacea Sacc. Syll. **8**: 770. 1889 (*fide* Massee).
Patinella olivacea Sacc. Syll. **8**: 770. 1889 (Batsch ?).
Patinella hirneola Sacc. Syll. **8**: 771. 1889 (*fide* Massee).
Patinella applanata Sacc. Syll. **8**: 771. 1889.
Humaria marchica Rehm, Rabenh. Kypt.-Fl. 1^3: 952. 1894.
Phaeopezia marchica Sacc. Syll. **11**: 415. 1895.
Aleurina marchica Sacc. & Syd. Syll. **16**: 739. 1902.
Humaria fuscocarpa Morgan, Jour. Myc. **8**: 189. 1902.
Aleurina fuscocarpa Sacc. & Syd. Syll. **16**: 739. 1902.
Catinella olivacea Boud. Hist. Class. Disc. Eur. 150. 1907 (Batsch ?).

Ascomata solitary or two or three together, sessile, attached to the substratum by numerous, radiating, dark brown fibers most

conspicuous in young plants; at first spherically closed, then opening out so as to become successively cup-shaped, saucer-shaped, finally applanate, with a permanently upturned margin; very young plants greenish yellow, becoming deeper and darker green, finally blackish olive when old, fleshy and somewhat gelatinous when fresh, 3–10 mm. in diameter, the majority 3–5 mm. On drying the disk becomes olive-black, the exterior brownish, somewhat furfuraceous or vertically striate, and the texture friable. *Excipulum* about 400 μ thick at the base, entirely parenchymatous, the ectal cells 18–20 μ in diameter, polygonal, with dark brown walls, becoming gradually smaller and yellow toward the hymenium, the superficial cells of the margin and sides projecting to form short, stout, obtuse, septate, flexuous, brown pili causing the surface to appear furfuraceous, those at the base much longer and radiating so as to form a small subiculum covering the substratum 2–3 mm. beyond the margin. On treatment with 1 per cent caustic potash the hymenium or whole flesh becomes deep violet and frequently yields a violet solution. *Hymenium* about 120–140 μ thick. *Asci* narrowly cylindric-clavate, apex rounded, not blue with iodine, 65–90 x 3–6 μ (mostly 70–80 μ). *Spores* eight, uniseriate, elliptic or elliptic-clavate, usually slightly narrowed near the middle so as to appear slipper-shaped, continuous, two-guttulate, pale olive-brown to deep brown, 7–11 x 4–5 μ (majority 8–10 μ); *Paraphyses* cylindric, septate, rarely branched, somewhat longer than the asci, the tips agglutinated with amorphous matter to form an epithecium.

On rotten wood, especially of old, moist, carious, decorticated logs, not abundant but widely distributed in eastern North America and the West Indies; also in Europe and Ceylon.

This is one of our most peculiar and attractive discomycetes. "In its young state it is truly *Peziza*-like, and very beautiful." While easily recognizable at sight when once understood, it has been described as new over and over again until very few fungi have a greater multiplicity of names. While this may be the species to which Batsch applied the name *Peziza olivacea*, his brief description and crude figures are much too inadequate to justify more than surmise, and may apply equally well to numerous other species. The apparent rarity of our plant in central Europe makes it still more improbable that Batsch had seen it. *P. olivacea* is simply one of the numerous names of fungi handed down from the pre-Persoonian period, the mycological stone age, which are absolutely indeterminable according to present taxonomic standards,

and which should therefore be dropped completely and no longer allowed to encumber the mycological literature.

The first certainly identifiable name applied to the plant under consideration is the one adopted in the present paper. While Schweinitz's type, from North Carolina, is missing from his herbarium, there is a specimen marked "rotten wood. Beth." under the name "*Lemalis pulla β nigro-olivascens* Schw. No. 1089." This label involves a curious switching of names and an error in spelling as follows: *Peziza nigro-olivacea* L. v. S. was included by Fries in his Systema as "*Patellaria pulla β nigro-olivacea.*" In 1834, Schweinitz transferred the species to the genus *Lemalis*, and changed the specific name so that it appeared as "1089. 4. *L. rufo-olivacea*, L. v. S., Syn. Car. 1220, *Peziza nigro-olivacea;* Salem et Bethlehem differt specifice a priori [*L. pulla* Fr.]". At a later date, Dr. Michener, in mounting and arranging Schweinitz's fungi, restored the original specific name (*nigro-olivacea*) to the label, but misspelled it "*nigro-olivascens.*" The presence of numbers and references in each case leaves no doubt that the specimen now in the Schweinitzian Herbarium is the one referred to in 1834 as from Bethlehem.

Although Fries stated that he had seen a specimen of *Peziza nigro-olivacea*, there seems to be no Schweinitzian material so labeled in the Friesian Herbarium, at Upsala. However, there is a specimen called "*Lemalis rufo-olivacea* Schw.," from Curtis, as well as one marked "*Peziza applanata* ex herb. Schweinitz." While both are typical *Catinella nigro-olivacea*, the latter is quite different from the specimen in Schweinitz's own herbarium called *P. applanata*.

In 1830, Fries described as *Bulgaria nigrita* a plant collected in Russia by Weinmann. So far as the writer can discover, no more recent collection has been referred to this species, which has remained practically unknown for nearly a century. In the Friesian Herbarium is a specimen marked "*Bulgaria nigrita*, Petersburg," in Elias Fries's own hand, which is in all probability the original type. When the writer saw it, in 1904, he was at once impressed with its gross resemblance to the American plant. Subsequent microscopical examination removed all doubts as to their specific identity.

During the Berkeleyan period this species was collected occasionally in various parts of the world, and almost as often described as new. Material from Cuba (Wright, No. 369) was given the name *Peziza viridi-atra* B. & C. Examination of the type, at Kew, shows it to be identical with the Schweinitzian plant. Saccardo placed it in *Pezicula*. Specimens from Ceylon were called *Patellaria applanata* B. & Br. The writer has not seen the type, but Massee, on examination, declared it to be *Patinella olivacea* (*C. nigro-olivacea*). On the basis of the same evidence the same writer included *P. violacea* B. & Br. and *P. hirneola* B. & Br., also from Ceylon, as synonyms. In the original description of *P. applanata* the authors mentioned a "United States specimen." Material so labeled in the herbaria of Fries and the Philadelphia Academy of Science, collected in South Carolina, by Curtis, has been examined, and certainly belongs here. Additional collections reported from Connecticut (Wright), Pennsylvania (Michener), and North Carolina (Curtis) have not been seen.

No specimen of *Rhizina nigro-olivacea* Curr. has been available to the writer. But since both Phillips and Massee (the latter after examination of the type) place it as a synonym of *P. olivacea*, there seems to be no doubt of its identity with *C. nigro-olivacea*.

Peziza fuscocarpa Ell. & Hol. was described from material collected in Iowa by Holway. It has been referred to *Phaeopezia* by Saccardo, to *Humaria* by Morgan, and to *Aleurina* by Saccardo & Sydow, and by Rehm. This specific name is the one under which the plant here considered has commonly been reported or distributed in America. The type is identical with *Peziza nigro-olivacea* L. v. S. The writer has received it from many localities and has had abundant opportunity to study it in the fresh condition in all stages of development.

Of all the references of the present species made by Saccardo, the most curious is that where *Peziza nigro-olivacea* L. v. S. is made a synonym of *Mollisia umbonata* (Pers.) Sacc. To Persoon's specific description are appended verbatim Fries's observations comparing Schweinitz's plant with *Patellaria pulla* Fr., a totally different plant from Persoon's. This confusion may be due to a mixing of notes.

Finally, the plant here considered was described as *Humaria marchica* by Rehm, from German material distributed by Sydow and examined by the writer.

While it has been demonstrated that *Catinella nigro-olivacea* has passed under about thirty different names, it may well be that others have escaped the writer's attention.

In the citation of specimens the following abbreviations have been used: CU, Herbarium of Cornell University; D, Durand Herbarium; F, Fries Herbarium; K, Kew Herbarium; Mo, Herbarium of the Missouri Botanical Garden; NY, Herbarium of the New York Botanical Garden; Phil, Philadelphia Academy; S, Schweinitz Herbarium.

MATERIAL EXAMINED

RUSSIA: Petersburg (F, type of *Bulgaria nigrita* Fr.).

GERMANY: Brandenburg, *Sydow* (NY, cotype of *Humaria marchica* Rehm, in Sydow, Myc. March. *2958*); Moravia, *Petrak* (D, as *Aleurina olivacea* [Batsch] v. Höhn.).

ONTARIO: Carleton Place, *Macoun* (NY); London, *Dearness* (CU, D, E. & E. N. Am. Fungi *2325*).

NEW HAMPSHIRE: Warren, *L. W. Riddle* (D).

NEW YORK: Ithaca, *Atkinson* (D); Varna, *Reddick* (D); Honeoye, *Durand* (D); Adirondack Mountains, Catskill Mountains, Old Forge, Kasoag, Big Indian, South Pond, Elisabethtown, *Peck* (A, as *Patellaria olivacea*).

PENNSYLVANIA: Bethlehem, *Schweinitz* (S); Springtown, *Witte* (NY); Mauch Chunk (Phil); Chester County, *Michener* (Phil).

NORTH CAROLINA: Blowing Rock, *Durand* (D).

SOUTH CAROLINA: *Curtis* (F, Phil).

LOUISIANA: Cloutierville & Abita Springs, *Langlois* (NY).

ARKANSAS: Camden, *C. J. Humphrey* (D).

OHIO: Preston, *Morgan* (NY).

MINNESOTA: Lakeville (D); St. Louis River, *Holway* (NY, Mo).

NORTH DAKOTA: Fargo, *Seaver* (NY, Mo, N. Dak. Fung. *28*, as *Phaeopezia fuscocarpa*).

IOWA: Mt. Pleasant, *Seaver* (D); Decorah, *Holway* (NY, type of *Peziza fuscocarpa*).

CUBA: *Wright 369* (K, type of *Peziza viridi-atra*)

2. **Catinella elastica** (Pat. & Gail.) Durand, comb. nov.

Phaeopezia elastica Pat. & Gail. Bull. Soc. Myc. Fr. **4**: 99. 1888.

Aleurina elastica (Pat. & Gail.) Sacc. & Syd. Syll. **16**: 739. 1902; Rehm, Ann. Myc. **1**: 515. 1903.

Ascomata sessile, solitary or in groups, 1–3 mm. in diameter, rich dark brown externally, black within, attached at the base by radiating brown fibers consisting of septate hyphae; margin thickened. *Excipulum* parenchymatous, of polygonal cells, the ectal ones with brown walls, those at the sides of the cup giving rise to short, septate, flexuous, brown pili, those at the base to stout, brown, radiating hyphae. *Asci* cylindric-clavate, apices rounded, not blue with iodine. *Spores* eight, uniseriate, elliptic, or elliptic-clavate, or slipper-shaped, continuous, brown, two-guttulate, 13–15 x 6–7 μ. *Paraphyses* cylindric, hyaline, septate, tips agglutinated with amorphous matter.

MATERIAL EXAMINED

VENEZUELA: on dead wood, Mapire, May, 1887, *A. Gaillard 6* (NY).

This species seems to differ from the preceding only in the larger spores. In the herbarium of the New York Botanical Garden is a specimen of what appears to be the original and only collection. The above description is that of the authors supplemented by notes drawn from the above mentioned specimen. In the original description the tissue is spoken of as elastic, somewhat like that of *Bulgaria*. The dried material is friable. It breaks up so that no complete ascus has been seen. The material becomes deep violet on treatment with KOH.

UNIVERSITY OF MINNESOTA

INDEX TO AMERICAN BOTANICAL LITERATURE

1908–1921

The aim of this Index is to include all current botanical literature written by Americans, published in America, or based upon American material; the word America being used in the broadest sense.

Reviews, and papers that relate exclusively to forestry, agriculture, horticulture, manufactured products of vegetable origin, or laboratory methods are not included, and no attempt is made to index the literature of bacteriology. An occasional exception is made in favor of some paper appearing in an American periodical which is devoted wholly to botany. Reprints are not mentioned unless they differ from the original in some important particular. If users of the Index will call the attention of the editor to errors or omissions, their kindness will be appreciated.

This Index is reprinted monthly on cards, and furnished in this form to subscribers at the rate of three cents for each card. Selection of cards is not permitted: each subscriber must take all cards published during the term of his subscription. Correspondence relating to the card issue should be addressed to the Treasurer of the Torrey Botanical Club.

Andres, H. Piroleen-Studien. Verh. Bot. Ver. Brandenburg **56**: 1–70. 6 Je 1914.

Arber, A. Leaves of the Helobieae. Bot. Gaz. **72**: 31–38. *pl. 1.* 16 Jl 1921.

Arthur, J. C. Specialization and fundamentals in botany. Am. Jour. Bot. **8**: 275–285. 30 Je 1921.

Bakhuizen Van den Brink, R. C. Revisio generis *Avicenniae*. Bull. Jard. Bot. Buitenzorg III. **3**: 199–226. *pl. 14–22.* My 1921.

Barnhart, J. H. Biographical notices of persons mentioned in the Schweinitz-Torrey correspondence. Mem. Torrey Club **16**: 290–300. 16 Jl 1921.

Blake, S. F. A new *Aspilia* from Trinidad. Proc. Biol. Soc. Washington **34**: 119, 120. 30 Je 1921.

Blake, S. F. New Meliaceae from Mexico. Proc. Biol. Soc. Washington **34**: 115–118. 30 Je 1921.

New species in *Cedrela* (1) and *Guarea* (2).

Bradshaw, R. V. Oregon willows. Am. Bot. **27**: 56–59. My 1921.

Britton, N. L. The cacti of Trinidad. Bull. Dept. Agr. Trinidad & Tobago **19**: 81–87. 1921.

Carano, E. Nuove ricerche sulla embriologia delle Asteraceae. Annali di Botanica **15**: 97–196. *pl. 4–12* + *f. 1, 2.* 28 F 1921.

Campbell, D. H. Professor H. Bruchmann. Science II. **54**: 67, 68. 22 Jl 1921.

Campbell, D. H. The eusporangiate ferns and the stelar theory. Am. Jour. Bot. **8**: 303–314. *f. 1–7.* 30 Je 1921.

Campbell, D. H. Die Verbreitung gewisser Lebermoose der malaiischen Region. Jahrb. Wissensch. Bot. **56**: 365–373. 1915.

Chamberlain, C. J. Helmut Bruchmann. Bot. Gaz. **72**: 45–47. 16 Jl 1921. [With portrait.]

Child, C. M. Certain aspects of the problem of physiological correlation. Am. Jour. Bot. **8**: 286–295. 30 Je 1921.

Clark, S. P. Sweet clover in Arizona. Univ. Arizona Coll. Agr. Circ. 34: 1–7. Ap 1921.

[Clute, W. N.] A new form of *Dentaria*. Am. Bot. **27**: 69. My 1921. *Dentaria laciniata* var. *dichronum* from Iowa.

Collins, J. L. Inbreeding and crossbreeding. Jour. Hered. **12**: 89–93. *f. 31–34.* 11 Jl 1921.

"The effect of inbreeding and crossbreeding in a wild plant [*Crepis capillaris*] of the sunflower family."

Conard, H. S. An outline for vascular plants. Science II. **54**: 15, 16. 1 Jl 1921.

Cook, M. T., & Martin, W. H. Potato diseases in New Jersey. New Jersey Agr. Exp. Sta. Circ. 122: 1–39. *f. 1–21.* F 1921.

Cook, M. T., & Poole, R. F. Diseases of sweet potatoes. New Jersey Agr. Exp. Sta. Circ. 123: 1–24. *f. 1–17.* Ap 1921.

Douin, C. La famille de Cephaloziellacées. Soc. Bot. France Mem. 29: 1–90. *f. 1–102.* 28 D 1920.

Includes as American novelties, the genus *Evansia*, and species in *Prionolobus* (1), *Evansia* (1), and *Cephaloziella* (9).

East, E. M., & Jones, D. F. Round tip tobacco—a plant "made to order." Jour. Hered. **12**: 51–56. *pl. 1* + *f. 1–5.* 11 Jl 1921.

Felt, E. P. Key to American insect galls. New York State Mus. Bull. 200: 1–310. *pl. 1–16* + *f. 1–250.* 1 Au 1917.

Fernald, M. L. The geographic distribution of hybrids. Science II. **54**: 73, 74. 22 Jl 1921.

Folsom, D. Potato leafroll. Maine Agr. Exp. Sta. Bull. 297: 37–52. *f. 26–35.* Ap 1921.

Fromme, F. D., & Wingard, S. A. Varietal susceptibility of beans to rust. Jour. Agr. Research **21**: 385–404. *pl. 69–73.* 15 Je 1921.

Glover, W. O. Blister canker of apple and its control. New York Agr. Exp. Sta. Bull. 485: 1–71. *pl. 1–15* + *f. 1–8.* Ja 1921.

Hansen, A. A. Protect our vanishing wild plants! Conservationist **4**: 67–70. My 1921. [Illust.]

Harms, H. Uber die Fluorescenz des Aufgusses der Rinde von *Prunus serotina*. Verh. Bot. Ver. Brandenburg **59**: 146–150. 15 Ja 1918.

Harms, H. Über Fluorescenz-Erscheinungen bei dem Holze der Leguminose *Eysenhardtia amorphoides* H. B. K. Verh. Bot. Ver. Brandenburg **56**: 184–197. 18 Ja 1915.

Harms, H. Nachträge und Verbesserungen zu meinem Aufsatze über Fluorescenzerscheinungen. Verh. Bot. Ver. Brandenburg **57**: 191–202. 1 Mr 1916.

Harms, H. Ernst Ule. Verh. Bot. Ver. Brandenburg **57**: 150–184. 1 Mr 1916. [With portrait.]

Harris, J. A. Tissue weight and water content in a tetracotyledonous mutant of *Phaseolus vulgaris*. Proc. Soc. Exper. Biol. and Med. **18**: 207–209. 1921.

Harvey, E. M., & Murneek, A. E. The relation of carbohydrates and nitrogen to the behavior of apple spurs. Oregon Agr. Coll. Exp. Sta. Bull. 176: 1–47. *f. 1–12.* Ap 1921.

Hayes, H. K., & Gardner, R. J. Breeding crop plants. Pp. i–xvii + 1–328. *f. 1–66.* New York. 1921.

Hume, A. M., Loomis, H., & Hutton, J. G. Water as a limiting factor in the growth of sweet clover (*M[elilotus] alba*). South Dakota Agr. Exp. Sta. Bull. 191: 257–298. *pl. 1, 2.* Jl 1920.

Hutchinson, C. B. Heritable characters of maize—VII. Shrunken endosperm. Jour. Hered. **12**: 76–83. *f. 20–24.* 11 Jl 1921.

Hutchinson, J. The genera of Fumariaceae and their distribution. Kew Bull. Misc. Inform. **1921**: 97–115. 1921.

Illick, J. S. The white pine. Am. For. **27**: 422–426. Jl 1921. [Illust.]

Jones, L. R., Walker, J. C., & Tisdale, W. B. *Fusarium* resistant cabbage. Univ. Wisconsin Agr. Exp. Sta. Research Bull. 48: 1–34. *f. 1–10.* N 1920.

Judd, C. S. The Alahee tree. Hawaiian For. & Agric. **18**: 133–137. Je 1921. [Illust.]

Kraemer, H. Plant colors. Am. Jour. Pharm. **93**: 414–416. Je 1921.

Kraemer, H. Some experiments on the modification of color in plants. Am. Jour. Pharm. **93**: 416–418. Je 1921.

Kunkel, L. O. A possible causative agent for the mosaic disease of corn. Bull. Exp. Sta. Hawaiian Sugar Pl. Assoc. **3**: 1–15. *pl. 4–15 + f. 1, 2.* 9 Jl 1921.

Lee, H. A. The relation of stocks to mottled leaf of *Citrus* trees. Philipp. Jour. Sci. **18**: 85–93. *pl. 1–3.* Ja 1921.

Loesener, T. Plantae Selerianae.—IX. Verh. Bot. Ver. Brandenburg **58**: 129–157. 1 My 1917.

Includes new American species in *Nemastylis* (1), *Iresine* (1), *Daphnopsis* (1), and *Aniseia* (1).

Lonay, H. Sur quélques genres rares ou critiques de Renonculacées. Bull. Soc. Bot. Belgique **45**: 191–204. 4 D 1908.

Long, W. H. Notes on new or rare species of rusts. Bot. Gaz. **72**: 39–44. 16 Jl 1921.

Includes new species in *Gymnosporangium* (1), and *Ravenelia* (3).

MacDougal, D. T. Water deficit and the action of vitamines, amino-compounds, and salts on hydration. Am. Jour. Bot. **8**: 296–302. 30 Je 1921.

Martin, W. H. Studies on tomato leaf-spot control. New Jersey Agr. Exp. Sta. Bull. 345: 1–42. *pl. 1 + f. 1.* N 1920.

Maxon, W. R. A neglected fern paper. Proc. Biol. Soc. Washington **34**: 111–114. 30 Je 1921.

Refers to Watt's Catalog of Ferns, published in Canadian Naturalist II. **13**: 157–160. 1867.

Merrill, E. D. New Philippine Moraceae. Philipp. Jour. Sci. **18**: 49–69. Ja 1921.

Merrill, E. D. New Philippine Myrtaceae. Philipp. Jour. Sci. **18**: 289–308. Mr 1921.

Mulford, F. L. Oaks for ornamental planting. Am. For. **27**: 461–467. Jl 1921. [Illust.]

Nash, G. V. *Acacia armata.* Addisonia **6**: 13. *pl. 199.* 29 Je 1921.

Nash, G. V. *Acacia leprosa.* Addisonia **6**: 7. *pl. 196.* 29 Je 1921.

Nash, G. V. *Acacia lineata.* Addisonia **6**: 15. *pl. 200.* 29 Je 1921.

Nash, G. V. *Acacia longifolia floribunda.* Addisonia **6**: 11. *pl. 198.* 29 Je 1921.

Nash, G. V. *Acacia Nabonnandi.* Addisonia **6**: 9. *pl. 197.* 29 Je 1921.

Nash, G. V. *Acacia pubescens.* Addisonia **6**: 3, 4. *pl. 194.* 29 Je 1921.

Nash, G. V. *Acacia pulchella.* Addisonia **6**: 5. *pl. 195.* 29 Je 1921.

Nash, G. V. *Acacia scapuliformis.* Addisonia **6**: 1, 2. *pl. 193.* 29 Je 1921.

Nelson, E. W. Lower California and its natural resources. Mem. Nat. Acad. Sci. **16**[1]: 1–171. *pl. 1–35.* 1921.

Neville, H. O. Pineapples in Cuba. Cuba Rev. **19**: 13–28. Jl 1921. [Illust.]

Osterhout, W. J. V., Thaxter, R., & Fernald, M. L. Lincoln Ware Riddle. Science II. **54**: 9. 1 Jl 1921.

Parish, S. B. A reminiscence of roses. Am. Bot. **27**: 60, 61. My 1921.

Patouillard, N. *Clathrotrichum*, nouveau genre d'hyphomycétes. Bull. Soc. Mycol. France **37**: 33–35. 15 Ap 1921.
Parasitic on *Setaria* in Ecuador.

Pittier, H. Acerca del genero *Gyranthera* Pittier. Bol. Com. e Indust. [Venezuela] **13**: 428–433. Ja 1921.
Includes *G. caribensis* sp. nov.

Pittier, H. Arboles y arbustos nuevos de Venezuela—I. Bol. Com. e Indust. [Venezuela] **13**: 417–428. Ja 1921.
Includes new species in *Inga* (1), *Raputia* (1), *Esenbeckia* (1), *Trichilia* (1), *Stillingia* (1), *Gustavia* (1), *Mouriria* (1), *Oxythece* (1), *Pouteria* (1), and *Sideroxylum* (1).

Rapp, C. W. Bacterial blight of beans. Oklahoma Agr. Exp. Sta. Bull. **131**: 1–39. *f. 1–17.* Jl 1920.

Reinking, O. A. Citrus diseases of the Philippines, southern China, Indo-China, and Siam. Philipp. Agric. **9**: 121–179. *pl. 1–14.* F 1921.

Reinking, O. A. Diseases of economic plants in Indo-China and Siam. Philipp. Agric. **9**: 181–183. F 1921.

Rolfe, R. A. *Wercklea lutea* [Malvaceae]. Kew Bull. Misc. Inform. **1921**: 118, 119. 1921.
New species from Costa Rica.

Rydberg, P. A. Two new species from Arizona. Am. Bot. **27**: 61–63. My 1921.
Eriogonum Clutei Rydb. and *Forestiera arizonica* (A. Gray) Rydb.

Sands, W. N. Plants common to the West Indies and Malaya. Agric. News **20**: 163; 182, 183. 28 My 1921.

Schmitz, H., & Daniels, A. S. Studies in wood decay—I. Laboratory tests on the relative durability of some western coniferous woods with particular reference to those growing in Idaho. School Forestry Univ. Idaho Bull. **1**: 1–11. Jl 1921.

Schmitz, H. Studies in wood decay—II. Enzyme action in *Polyporus volvatus* Peck and *Fomes ignarius* (L.) Gillet. Jour. Gen. Physiol. **3**: 795–800. 20 Jl 1921.

Shear, C. L., & Stevens, N. E. (Editors). The correspondence of Schweinitz and Torrey. Mem. Torrey Club **16**: 119–300. *pl. 6, 7.* 16 Jl 1921.

Sherman, H. Respiration of dormant seeds. Bot. Gaz. **72**: 1–30. *f. 1–4.* 16 Jl 1921.

Soth, M. E. Goat-weed in Idaho. Am. Bot. **27**: 70. My 1921.

Sousa Brito, E. C. de. Algumas plantas uteis do Brasil, o hybridismo de uma dellas. Arch. Esc. Sup. Agric. e Med. Vet. **4**: 25–54. *pl. 10–16.* D 1920.

Stadler, L. J., & Helm, C. A. Corn in Missouri—I. Corn varieties and their improvement. Missouri Agr. Exp. Sta. Bull. 181: 1–51. *f. 1–7.* Mr 1921.

Stahel, G. De Sclerotium-ziekte van de Liberiakoffie in Suriname veroorzaakt door *Sclerotium coffeicolum* nov. spec. Dept. Landb. Suriname Bull. 42: 1–34. *pl. 1–11.* Ja 1921.

Stevens, F. L. The relation of plant pathology to human welfare. Am. Jour. Bot. **8**: 315–322. 30 Je 1921.

Stout, A. B. Types of flowers and intersexes in grapes with reference to fruit development. New York Agr. Exp. Sta. Techn. Bull. 82: 1–16. *pl. 1–7.* Ja 1921.

Sydow, H. & P. Novae fungorum species—XVI. Ann. Mycol. **18**: 154–160. Ap 1921.

Includes new American species in *Septobasidium* (1), *Uromyces* (1), *Aecidium* (1), *Catacauma* (1), *Taphrina* (1).

Taubenhaus, J. J. A study of the black and the yellow molds of ear corn. Bull. Texas. Agr. Exp. Sta. **270**: 3–38. *f. 1–10.* O 1920.

Taubenhaus, J. J., & Mally, F. W. Pink root disease of onions and its control in Texas. Texas Agr. Exp. Sta. Bull. 273: 1–42. *f. 1–3.* Ja 1921.

Taylor, W. R. Additions to the flora of Mount Desert, Maine. Rhodora **23**: 65–68. 20 Ap 1921.

Thatcher, R. W. The chemistry of plant life. Pp. i–xvi + 1–268. New York. 1921.

Thomson, G. E. Hegari in Arizona. Univ. Arizona Coll. Agr. Circ. 33: 1–4. Ap 1921. [Illust.]

Thomas, R. C. *Botrytis* rot and wilt of tomato. Bull. Ohio Agr. Exp. Sta. 6: 59–62. Ap 1921. [Illust.]

Thurston, H. W. A note on the corrosive sublimate treatment for the control of *Rhizoctonia*. Phytopathology 11: 150, 151. Mr 1921.

Tiffany, L. H. Algal food of the young gizzard shad. Ohio Jour. Sci. 21: 113–122. F 1921.

Tiffany, L. H. New forms of *Oedogonium*. Ohio Jour. Sci. 21: 272, 273. *pl. 1*. Je 1921.
2 new species from Ohio.

Tisdale, W. H., & Jenkins, J. M. Straighthead of rice and its control. U. S. Dept. Agr. Farm. Bull. 1212: 1–16. *f. 1–6*. Je 1921.

Toumey, J. W. Damage to forests and other vegetation by smoke, ash and fumes from manufacturing plants in Naugatuck Valley, Connecticut. Jour. For. 19: 367–373. *f. 1–6*. Ap 1921.

Trelease, W. The Jack Oak (*Quercus ellipsoidalis*). Trans. Illinois Acad. Sci. 12: 108–118. *pl. 139–143*. 1919.

Trelease, W. North American pipers of the section *Ottonia*. Am. Jour. Bot. 8: 212–217. *pl. 5–8*. 30 Ap 1921.

Tucker, E. M. Bibliographical notes. Jour. Arnold Arbor. 2: 181–184. Ja 1921.
Discussion of issuance of Nouveau Duhamel and of Torrey and Gray's Flora of North America.

Tucker, E. M. Catalogue of the library of the Arnold Arboretum of Harvard University. Publ. Arnold Arbor. 6. 1: 1–782. 1914; 2: 1–542. 1917.
Compiled under the direction of Charles Sprague Sargent.

Ule, E. Unter den Indianern des nördlichen Amazonas. Verh. Naturw. Vereins Hamburg III. 21: LXXVI, LXXVII. 1914.
A botanical and anthropological journey.

Uphof, J. C. T. Die Waldflora im Staate Washington. Vegationsbilder XIII. 7: *11 pp. + pl. 37–42*. 1920.

Urban, I. Novitates haitienses. Notizbl. Bot. Gart. und Mus. Berlin-Dahlem 7: 495–499. 1 Mr 1921.
Includes new species in *Pilea* (1), *Phthirusa* (1), *Cassia* (1), *Sebastiana* (1). *Abutilon* (1) and *Eugenia* (1).

Urban, I. Über die Malvaceen-Gattung *Montezuma* Moc. et Sessé. Notizbl. Bot. Gart. und Mus. Berlin-Dahlem **7**: 543, 544. 1 Mr 1921.

Wakefield, E. M. Mosaic diseases of plants. West Indian Bull. **18**: 197–206. [1921.]

Walker, E. R. The gametophytes of *Equisetum laevigatum*. Bot. Gaz. **71**: 378–391. *pl. 23, 24 + f. 1–3*. 16 My 1921.

Walker, J. C. Onion smudge. Jour. Agr. Research **20**: 685–721. *pl. 80–85 + f. 1–10*. 1 F 1921.

Walker, J. C. Rust of onion followed by a secondary parasite. Phytopathology **11**: 87–90. *f. 1, 2*. F 1921.

Weatherby, C. A. A form of *Ilex opaca*. Rhodora **23**: 118, 119. 19 My 1921.

Weir, J. R. *Thelephora terrestris*, *T. fimbriata*, and *T. caryophyllea* on forest tree seedlings. Phytopathology **11**: 141–144. *pl. 5*. Mr 1921.

Wells, B. W. Evolution of zoocecidia. Bot. Gaz. **71**: 358–377. *pl. 21, 22*. 16 My 1921.

Wells, B. W. A phenomenal shoot. Science II. **54**: 13, 14. 1 Jl 1921.
Paulownia tomentosa.

Weston, W. H. Another conidial *Sclerospora* of Philippine maize. Jour. Agr. Research **20**: 669–684. *pl. 76–79*. 1 F 1921.

Whaley, W. Properties of wood. A chapter from Lucien Chancerel's Botanique Forestière. Jour. For. **19**: 414–420. Ap 1921.

White, O. E. The pollination of flowers. Brooklyn Bot. Gard. Leafl. **9** n. 3, 4: 1–15. 4 My 1921.

Wiegand, K. M. *Amelanchier amabilis*, a new name. Rhodora **23**: 48. 5 Ap 1921.

Wiegand, K. M. *Echinochloa* in North America. Rhodora **23**: 49–65. 20 Ap 1921.
Includes *E. paludigena* sp. nov. and several new varieties.

Wieland, G. R. Monocarpy and pseudomonocarpy in the cycadeoids. Am. Jour. Bot. **8**: 218–230. *pl. 9–12*. 30 Ap 1921.

Williams, C. B. Report on the Froghopper Blight of sugar-cane in Trinidad. Mem. Dept. Agr. Trinidad and Tobago **1**: 1–170. *pl. 1–11 + f. 1–32*. Ja 1921.
Considers also "The Root Disease of Sugar Cane" caused by fungi.

Yunker, T. G. A Curious abnormality in *Cuscula cvspidata*. Am. Bot. **27**: 48, 49. My 1921. [Illust.]



1 2

PESSIN : EPIPHYLLOUS PLANTS OF JAMAICA